



MALARIA EPIDEMIC EARLY WARNING PREDICTION SYSTEM FOR WESTERN KENYA HIGHLAND FOR MARCH 2025

Ref No: MET/7/58/03-2025

Issue Date: 03/03/2025

1. Summary

The model outputs for the malaria epidemic early prediction system for the western highlands of Kenya indicate no risk of Malaria in all the three areas in the months of March, 2025 and April, 2025

2. Model Outputs

2.1 Malaria epidemic early prediction system for Kakamega

Table 1 below shows the malaria epidemic early prediction system for Kakamega for March, 2025.

Table 1: MALARIA EPIDEMIC EARLY PREDICTION SYSTEM: KAKAMEGA


Yr.	Month	Tmax	Mean Tmax	Tmax Deviation /anomaly	R/fall (mm)	R/fall Code	Tmax Deviation /anomaly Code	Additive % Risk
2025	1	29.9	28.3	1.6	85.3	0	4	18.2
2025	2	32.7	29.2	3.5	3.9	0	16	18.2

The observed climate data for February, 2025 indicates an increase in maximum temperature from 29.9°C in January 2025 to 32.7°C in February, 2025. This observation in January, 2025 *was positive (3.5 above the mean of the month)*. Rainfall decreased from 85.3mm in January, 2025 to 3.9mm in February, 2025. The additive model percentage risk is **18.2%**.

Box 1:
For Kakamega, the epidemic threshold level is **30%**.

Consequently, there is no risk of Malaria Epidemic in Kakamega in the month of March, and April, 2025 (See Figure 1)

Figure 1:



2.2 Malaria epidemic early prediction system for Kisii

Table 2 below shows the malaria epidemic early prediction system for Kisii for March, 2025.

Table 2: MALARIA EPIDEMIC EARLY PREDICTION SYSTEM: KISII

Yr	Mon	Tmax (°C)	Mean Tmax (°C)	Tmin (°C)	Mean Tmin (°C)	Tmax Dev./anom	Tmin Dev./anom	Total Temp Dev./Anom	Temp Dev./anom Code	R/fall (mm)	R/fall Code	Model Output
2025	1	29.9	26.1	16.9	15.7	3.8	1.2	5.0	5	206.4	1	0
2025	2	29.2	27.0	11.3	16.1	2.2	-4.8	-2.6	0	47.6	0	0

The observed climate data for Kisii for February, 2025 indicates a decrease in maximum temperature from 29.9°C in January, 2025 to 29.2°C in February, 2025.

This observation in February, 2025 was *positive (2.2 above the mean of the month)*. Rainfall decreased from 206.4mm in January, 2025 to 47.6mm in February, 2025.

Box 2:
For Kisii, the epidemic threshold level is 20%.

The model output risk is **Nil**. Therefore, there is no risk of malaria epidemic in Kisii in the month of March, and April, 2025. (See Figure 2).

Figure 2:




Figure 2: Time series of the risk of malaria outbreak in Kisii

2.3 Malaria epidemic early prediction system for Nandi

Table 3 below shows the malaria epidemic early prediction system for Nandi for March, 2025.

Table 3: NANDI MALARIA EPIDEMIC EARLY PREDICTION SYSTEM

Yr	M0n	Tmax (°C)	Mean Tmax (°C)	Tmax Dev.	Tmin	Mean Tmin	Tmin Dev. /anom	Total Temp Dev. /Anom	R/fall (mm)	Temp Dev. Filters	R/fall Filters	Multiplicative Model
2025	1	25.1	23.3	1.8	11.8	10.9	0.9	2.7	101	3	0	0
2025	2	27.6	23.2	4.4	16.8	11.7	5.1	9.5	32.7	5	0	0


The maximum temperature in Nandi indicates an increase from 25.1°C in January, 2025 to 27.6°C in February, 2025. This observation in February, 2025 for Nandi was *positive (4.4°C above the mean of the month)*. Rainfall decreased from 101mm in January, 2025 to 32.7mm in February, 2025.

Box 3:
For Nandi, epidemic threshold level is **20%**.

The additive model percentage risk is **Nil**.

Hence, there is no expectancy for a malaria outbreak for the month of March, and April, 2025. (See Figure 3)

Figure 3:



3. Disclaimer

The information presented in this bulletin is based on [predictive models and observed climate data](#), which are subject to change. While every effort has been made to ensure the accuracy and reliability of the data, the following points should be noted.

Public Health Advisory: This bulletin is intended for informational purposes only. It should not be used as the sole basis for public health decisions. Local health authorities should be consulted for actionable guidance and preventive measures against malaria.

Continuous Monitoring: Malaria transmission dynamics are influenced by numerous factors, including temperature, rainfall, and human behaviour. Continuous monitoring and updates to the predictive models are essential for accurate assessments.

Updates: This bulletin reflects data and predictions as of March 2025. Future updates will be issued as new data becomes available.

Dr Gikungu

DIRECTOR, KENYA METEOROLOGICAL DEPARTMENT